

A Large-D Weyl Invariant String Model in Anti-de Sitter Space

Ian Davies* and Paul Mansfield†

*Centre for Particle Theory, Department of Mathematical Sciences
University of Durham, South Road, Durham DH1 3LE, U.K.*

Abstract

Strings propagating in AdS_{D+1} are made Weyl invariant to leading order in large D by a ghost-matter coupling that preserves the Poincaré symmetry of the boundary.

It has been proposed [1] [2] that Wilson loops for gauge theories in flat space with coordinates X^i are described by strings in a target space with metric of the form

$$ds^2 = d\varphi^2 + z^2(\varphi) \sum_{i=1}^D (dX^i)^2. \quad (1)$$

The extra coordinate φ is the Liouville mode and the ‘warp factor’ is taken to be

$$z^2(\varphi) = \exp\left(\frac{2\varphi}{l}\right)$$

so that the metric is that of Euclidean AdS_{D+1} space with radius of curvature l . (This metric also occurs in brane-world models and in the AdS/CFT correspondence.) When the curvature is small on the string scale, i.e. when $\alpha'/l^2 \ll 1$, the conditions for Weyl invariance of the string theory can be reliably computed in perturbation theory and require the presence of X^i -dependent fields such as the dilaton [3]. These break the space-time symmetries of the original flat space and add to the complexity of the model. However one would also like to investigate the regime where α'/l^2 cannot be assumed small so that the finite length of the string plays a significant role. Such situations can arise when the curvature of the target space geometry is appreciable at the string scale. In order to probe such situations one must either develop non-perturbative techniques, or identify some other dimensionless parameter in the theory which can be taken to be small in the regime of interest. One could then perform a perturbative computation of the Weyl anomaly with such a parameter and see how the conventional conditions for Weyl invariance are changed. We will analyze string

*I.J.Davies@durham.ac.uk

†P.R.W.Mansfield@durham.ac.uk

propagation in the target space of the form (1) in an expansion in inverse powers of the dimension. D plays an analogous role to N in the $O(N)$ nonlinear sigma model, which has a well-known large N expansion [4]. By working to leading order we obtain a Weyl invariant theory by tuning D such that $D+1 = 26$, and by including a coupling between the Faddeev-Popov ghost sector of the worldsheet theory and the “matter” field φ . Such a coupling can, in part, be interpreted as the presence of a dilaton field that depends only on φ . In the context of Polyakov’s conjecture [5], this means that the dilaton no longer breaks Poincaré and rotational invariance in the boundary gauge theory, unlike the dilaton field that one requires in the conventional small α'/l^2 Weyl anomaly analysis. (Coupling matter and ghost sectors has been studied in different contexts in [6] [7].)

To pursue the analogy with the $O(N)$ nonlinear sigma model we write the string partition function in terms of coordinates $W^i = zX^i$ so that

$$Z = \int \mathcal{D}g \mathcal{D}z \mathcal{D}W \exp(-S[g, W, z])$$

with

$$S[g, W, z] = \frac{1}{4\pi} \int d^2\xi \sqrt{g} \left[g^{ab} l^2 \frac{\partial_a z \partial_b z}{z^2} + W^i \left(\Delta - \frac{1}{z} \Delta z \right) W^i \right]$$

Δ is the usual covariant worldsheet Laplacian, and we work in units such that α' is equal to one. The indices a, b label the two worldsheet coordinates ξ_1, ξ_2 . We ignore world sheet boundary effects. Performing the Gaussian W -integral generates the functional determinant of the operator

$$\Gamma = \left(\Delta - \frac{1}{z} \Delta z \right)$$

The explicit dependence of $\text{Det}^{-D/2}(\Gamma)$ on the metric scale $\phi(\xi)$ in the conformal gauge $g_{ab}(\xi) = e^{\phi(\xi)} \delta_{ab}$ can be found using heat kernel techniques,

$$\begin{aligned} \text{Det}^{-D/2}(\Gamma) &= \exp \left(\int d^2\xi \left[\frac{D}{96\pi} (\partial_a \phi)^2 - \frac{D}{8\pi} \frac{\phi}{z} \partial_a^2 z + \dots \right] \right) \\ &\equiv \exp \left(\int d^2\xi [A + B + \dots] \right) \end{aligned} \quad (2)$$

where $(\dots)$ represents ϕ -independent terms that we will consider in a moment. The first term in (2), A , is the usual contribution to the Weyl anomaly from D free bosons [8], and can be combined with the standard Faddeev-Popov gauge fixing determinant to give a factor of

$$\exp \left(\frac{D-26}{96\pi} \int d^2\xi [(\partial_a \phi)^2 + \lambda e^\phi] \right) = \exp \left(\frac{D-26}{96\pi} S_L \right)$$

where S_L is the Liouville action for $\phi(\xi)$ in conformal gauge. (The integral over z will produce a further contribution to this.)

In critical string theory the overall ϕ dependence of Z must be cancelled for consistency. This is also true in non-critical string theory since the split of the scale

of the metric into the dynamical φ and the fiducial ϕ is arbitrary [9] [10]. Clearly terms proportional to S_L can be cancelled by tuning D (to a large value).

The second term in (2), B , is more problematic. In the usual small α'/l^2 calculation similar terms would appear from expanding the action in Gaussian normal coordinates about a point, so

$$G_{\mu\nu}(\hat{X})\partial_a\hat{X}^\mu\partial_b\hat{X}^\nu = \delta_{\mu\nu}\partial_ax^\mu\partial_bx^\nu - \frac{1}{3}R_{\mu\lambda\nu\kappa}(C)x^\lambda x^\kappa\partial_ax^\mu\partial_bx^\nu + \dots$$

where $\hat{X}^\mu = C^\mu + x^\mu$. Contracting x^λ with x^κ produces a term proportional to $\phi R_{\mu\nu}\partial_ax^\mu\partial_bx^\nu$. A partial integration would turn this into something resembling B when the latter is expanded about a constant value of z . Further contraction of ∂_ax^μ with ∂_bx^ν will contribute to the dilaton beta function at sub-leading order. The term in (2) is, by contrast, of leading order in D . As we will see the z -propagator is different at leading order in the large- D expansion, consequently B does not contribute to the dilaton beta-function, but it does make the z -propagator depend on ϕ . This does not happen in the small α'/l^2 calculation because $x\partial^2x$ is proportional to the equation of motion. So to make the z -propagator independent of ϕ at leading order in D we need to cancel B . This can be done without breaking the symmetries of the X space if we couple z to the ghosts.

Firstly “bosonize” the Faddeev-Popov ghosts by writing the Faddeev-Popov determinant as

$$\text{Det}'_{FP} = \exp\left(\frac{26}{96\pi}S_L\right) \sim \int \mathcal{D}\psi \exp\left(-\frac{1}{96\pi} \int d^2\xi \sqrt{g} [\psi\Delta\psi + 2\beta R^{(2)}\psi]\right)$$

where $R^{(2)}$ is the scalar curvature of g_{ab} , the prime denotes omission of the zero modes, and $\beta^2 = -25$. Adding

$$S_{ct} = \frac{D}{8\pi\beta} \int d^2\xi \sqrt{g} \frac{\psi}{z} \Delta z$$

to the action cancels B in equation (2) on integration over $\psi(\xi)$. Integration by parts shows that this counterterm can be interpreted as a dilaton field [11] of the form $\Phi(z) \sim \ln(z)$, as well as a coupling of the target space metric to the ghost $\psi(\xi)$ of the form $ds_{gh}^2 \sim (\psi/z)dz^2$.

In order to complete the calculation of the partition function we must also determine the z -dependence of $\text{Det}^{-D/2}(\Gamma)$. This is facilitated by the $1/D$ expansion. We write the $z(\xi)$ field as

$$z(\xi) = z_0 + \frac{\bar{z}(\xi)}{\sqrt{D}}$$

where z_0 is a constant on the world-sheet and $\bar{z}(\xi)$ are quantum fluctuations about z_0 to be integrated over. To obtain an effective action for the $\bar{z}(\xi)$ fluctuations, the determinant of Γ is expanded in powers of $\bar{z}(\xi)$ retaining terms up to $O(1)$ in D . This is most easily done in momentum space and gives the leading order effective theory

$$Z \sim \exp\left(\frac{D-26}{96\pi}S_L\right) \times \int \mathcal{D}\bar{z} \exp\left(\frac{1}{8\pi} \int \frac{d^2p}{(2\pi)^2} \frac{1}{z_0^2} \bar{z}(p)\bar{z}(-p)p^2 \ln\left(\frac{p^2}{\Lambda^2}\right)\right) \quad (3)$$

(note that the Fourier transform of the $\bar{z}$ -propagator is not the usual $1/p^2$, as was mentioned earlier. It is the improved ultra-violet behaviour that is responsible for B not contributing to the dilaton beta-function in next to leading order). Λ^2 is a finite overall scale. We need to integrate over the fluctuations to obtain the final form of the Weyl anomaly, and this will involve computing the ϕ -dependence of the determinant of the operator whose Fourier representation is given in equation (3). Again, heat kernel techniques and the fact that this operator derives from Γ (whose ϕ -dependence is readily found to be $\delta_\phi \Gamma = -\delta\phi \Gamma$) gives a contribution to the Weyl anomaly equal to that of a single free boson. Hence, the final Weyl anomaly is found to be

$$Z \sim \exp \left(\frac{D+1-26}{96\pi} S_L \right)$$

and so Z is Weyl invariant when $D+1=26$, as usual.

In conclusion we have found that to leading order in a large- D expansion it is possible to construct a Weyl invariant string theory in Euclidean AdS_{D+1} space without breaking the symmetries associated with the boundary. This is achieved by coupling the z coordinate to the ghosts, resulting in part in a z -dependent dilaton. The $1/D$ expansion promotes to leading order terms that are usually negligible for small α'/l^2 . The question of whether the ghost-matter coupling is necessary to achieve Weyl invariance, rather than just sufficient, is one which warrants further investigation, as is the question of next to leading order effects.

Acknowledgements

Ian Davies would like to thank James Gregory, Ken Lovis, Tony Padilla and David Page for helpful discussions. Ian Davies is supported by a PPARC research studentship.

References

- [1] A. M. Polyakov, “String theory and quark confinement”, Nucl. Phys. Proc. Suppl. **68** (1998) 1, [hep-th/9711002](#).
- [2] S. S. Gubser, I. R. Klebanov, and A. M. Polyakov, “Gauge theory correlators from non-critical string theory”, Phys. Lett. **B428** (1998) 105, [hep-th/9802109](#).
- [3] C. G. Callan, E. J. Martinec, M. J. Perry, and D. Friedan, “Strings in background fields”, Nucl. Phys. **B262** (1985) 593.
- [4] A. M. Polyakov, “Gauge fields and strings”, Chur, Switzerland: Harwood (1987) 301p. (Contemporary Concepts in Physics, 3).
- [5] A. Polyakov and V. Rychkov, “Gauge fields - strings duality and the loop equation”, Nucl. Phys. **B581** (2000) 116, [hep-th/0002106](#).

- [6] E. Witten, “Chern-Simons gauge theory as a string theory”, [hep-th/9207094](#).
- [7] W. Siegel, “Covariantly second-quantized strings 1,2”, *Phys. Lett.* **B151** (1985) 391,396.
- [8] A. M. Polyakov, “Quantum geometry of bosonic strings”, *Phys. Lett.* **B103** (1981) 207.
- [9] F. David, “Conformal field theories coupled to 2-D gravity in the conformal gauge”, *Mod. Phys. Lett.* **A3** (1988) 1651.
- [10] J. Distler and H. Kawai, “Conformal field theory and 2-D quantum gravity or Who’s afraid of Joseph Liouville?”, *Nucl. Phys.* **B321** (1989) 509.
- [11] T. Banks, D. Nemeschansky, and A. Sen, “Dilaton coupling and BRST quantization of bosonic strings”, *Nucl. Phys.* **B277** (1986) 67.